

Home Health ICD-9/ICD-10 Alert

You Be the Coder: Check OASIS Responses with DTI

Question: Our new patient fell at home and he remained on the floor for three days until his daughter found him. He has a large purplish bruise over his sacral area. Two days later he developed a large necrotic area over the sacrum and it is documented as a suspected deep tissue injury (DTI). We have orders to provide skilled nursing to assess integumentary status. How should we code for him?

Connecticut Subscriber

Answer: Code for this patient as follows, says **Joan L. Usher, BS, RHIA, COS-C, ACE**, with **JLU Health Record Systems** in Pembroke, Mass.:

- **M1020:** 707.03 (Pressure ulcer; lower back);
- **M1022b:** 707.25 (Pressure ulcer; unstageable); and
- **M1022c:** V15.88 (At risk for falling).

A DTI is a purple or maroon localized area of discolored intact skin or blood filled blister due to damage of underlying soft tissue from pressure and/or shear, Usher says. Code for these wounds as unstageable pressure ulcers, and be sure they are also reported accurately in the relevant OASIS items.

For M1306 Does this patient have at least one Unhealed Pressure Ulcer at Stage II or Higher or designated as unstageable the correct response for this patient is "Yes."

In M1308 Current number of unhealed [non-epithelialized] pressure ulcers, you'll count this wound in d.3 Unstageable: Suspected deep tissue injury in evolution.

Your M1320 Status of most problematic pressure ulcer is "Non-healing" because DTIs do not granulate and are not covered with new epithelial tissue.